Daftar Pustaka

[1] Emmanuel Audusse, François Bouchut, Marie-Odile Bristeau, Rupert Klein, and Benoı t Perthame. A fast and stable well-balanced scheme with hydrostatic reconstruction for shallow water flows. *SIAM Journal on Scientific Computing*, 25(6):2050–2065, 2004.

[2] Emmanuel Audusse, Marie-Odile Bristeau, and Benoit Perthame. *Kinetic schemes for Saint-Venant equations with source terms on unstructured grids*. PhD thesis, INRIA, 2000.

[3] Francesco Bassi and Stefano Rebay. High-order accurate discontinuous finite element solution of the 2d euler equations. *Journal of computational physics*, 138(2):251–285, 1997.

[4] Clothilde Bon. *Modélisation et simulation numérique d'écoulements hydrauliques et de ruissellement en topographie quelconque*. PhD thesis, Bordeaux 1, 1997.

[5] N Botta, R Klein, S Langenberg, and S Lützenkirchen. Well balanced finite volume methods for nearly hydrostatic flows. *Journal of Computational Physics*, 196(2):539–565, 2004.

[6] François Bouchut. Nonlinear stability of finite volume methods for hyperbolic conservation laws. 2004.

[7] A Chinnayya and AY LeRoux. A new general riemann solver for the shallow-water equations with friction and topography, 1999. *Available in the conservation law preprint server http://www. math. ntnu. no/conservation*.

[8] Taichung Fong. Two-dimensional numerical modelling of hyperconcentrated flows.

[9] Thierry Gallouët, Jean-Marc Hérard, and Nicolas Seguin. Some approximate godunov schemes to compute shallow-water equations with topography. *Computers & Fluids*, 32(4):479–513, 2003.

[10] N Goutal and F Maurel. *Proceedings of the 2nd workshop on dam-break wave simulation*. Electricité de France. Direction des études et recherches, 1997.

[11] Peng Hu, Zhixian Cao, and Gareth Pender. Well-balanced two-dimensional coupled modelling of submarine turbidity currents. In *Proceedings of the Institution of Civil Engineers-Maritime Engineering*, volume 165, pages 169–188. Thomas Telford Ltd, 2012.

[12] Rupert Klein, N Botta, and AZ Owinoh. Distinguished limits, multiple scales asymptotics, and numerics for atmospheric flows. 2001.

[13] AY LeRoux. Discrétisation des termes sources raides dans les problemes hyperboliques. *Systemes hyperboliques: noveau schemas et nouvelles applications. Ecoles CEA-EDFINRIA "problemes non linéaires appliqués", INRIA Rocquencourt (France)*, 1998.

[14] Sudi Mungkasi. Metode volume hingga untuk menyelesaikan masalah bendungan-bobol. *Jurnal Mat Stat*, 11(01), 2011.

[15] James J Quirk. A contribution to the great riemann solver debate. In *Upwind and High-Resolution Schemes*, pages 550–569. Springer, 1997.