

DAFTAR PUSTAKA

- [1] K. Lehnertz, “Epilepsy and nonlinear dynamics,” *J. Biol. Phys.*, vol. 34, no. 3-4 SPEC. ISS., pp. 253–266, 2008.
- [2] T. Tomson, L. Nashef, and P. Ryvlin, “Sudden unexpected death in epilepsy: current knowledge and future directions,” *Lancet Neurol.*, vol. 7, no. 11, pp. 1021–1031, 2008.
- [3] “Epilepsy,” 2019. [Online]. Available: <https://www.who.int/news-room/fact-sheets/detail/epilepsy>. [Accessed: 21-Oct-2019].
- [4] S. L. Moshé, E. Perucca, P. Ryvlin, and T. Tomson, “Epilepsy: New advances,” *Lancet*, vol. 385, no. 9971, pp. 884–898, 2015.
- [5] G. Gerard, D. Shabas, and D. Rossi, “MRI in epilepsy,” *Comput. Radiol.*, vol. 11, no. 5–6, pp. 223–227, 1987.
- [6] M. R. Sperling, G. Wilson, J. Engel, T. L. Babb, M. Phelps, and W. Bradley, “Magnetic resonance imaging in intractable partial epilepsy: Correlative studies,” *Ann. Neurol.*, vol. 20, no. 1, pp. 57–62, 1986.
- [7] D. K. Binder and S. R. Haut, “Toward new paradigms of *seizure* detection,” *Epilepsy Behav.*, vol. 26, no. 3, pp. 247–252, 2013.
- [8] K. Doi, “Computer-aided diagnosis in medical imaging: Historical review, current status and future potential,” *Comput. Med. Imaging Graph.*, vol. 31, no. 4–5, pp. 198–211, 2007.
- [9] I. Wijayanto and A. Rizal, “Epileptic *Seizure* Detection in EEG Signal using EMD and Entropy,” *J. Electron. Syst.*, vol. 9, no. 2, p. 44, 2019.
- [10] Q. Yuan, W. Zhou, S. Yuan, X. Li, J. Wang, and G. Jia, “Epileptic EEG classification based on kernel sparse representation,” *Int. J. Neural Syst.*, vol. 24, no. 4, pp. 1–13, 2014.
- [11] A. Shoeb, “Application of machine learning to epileptic *seizure* onset detection and treatment,” *Diss. Massachusetts Inst. Technol.*, pp. 157–162, 2009.
- [12] N. Kannathal, M. L. Choo, U. R. Acharya, and P. K. Sadasivan, “Entropies for detection of epilepsy in EEG,” *Comput. Methods Programs Biomed.*, vol. 80, no. 3, pp. 187–194,

2005.

- [13] R. W. Homan, J. Herman, and P. Purdy, “Cerebral location of international 10-20 system electrode placement,” *Electroencephalogr. Clin. Neurophysiol.*, vol. 66, no. 4, pp. 376–382, 1987.
- [14] H. Fauzi, M. A. Azzam, M. I. Shapiai, M. Kyoso, U. Khairuddin, and T. Komura, “Energy extraction method for EEG channel selection,” *Telkomnika (Telecommunication Comput. Electron. Control.)*, vol. 17, no. 5, pp. 2561–2571, 2019.
- [15] H. Fauzi, M. I. Shapiai, S. Shah Abdullah, and Z. Ibrahim, “Automatic Energy Extraction Methods for EEG Channel Selection,” *Proc. - 2018 Int. Conf. Control. Electron. Renew. Energy Commun. ICCEREC 2018*, pp. 70–75, 2018.
- [16] “12. Signal Energy and Power 12.1.,” pp. 0–5.
- [17] N. E. Huang *et al.*, “The empirical mode decomposition and the Hubert spectrum for nonlinear and non-stationary time series analysis,” *Proc. R. Soc. A Math. Phys. Eng. Sci.*, vol. 454, no. 1971, pp. 903–995, 1998.
- [18] R. Esteller *et al.*, “Fractal dimension characterizes *seizure* onset in epileptic patients,” pp. 2343–2346 vol.4, 2008.
- [19] B. P. Harne, “Higuchi fractal dimension analysis of EEG signal before and after OM chanting to observe overall effect on brain,” *Int. J. Electr. Comput. Eng.*, vol. 4, no. 4, pp. 585–592, 2014.
- [20] C. Sevcik, “On fractal dimension of waveforms,” *Chaos, Solitons and Fractals*, vol. 28, no. 2, pp. 579–580, 2006.
- [21] B. S. Raghavendra and D. N. Dutt, “Computing fractal dimension of signals using multiresolution box-counting method,” *World Acad. Sci. Eng. Technol.*, vol. 37, pp. 1266–1281, 2010.
- [22] I. Wijayanto, R. Hartanto, H. A. Nugroho, and B. Winduratna, “Seizure Type Detection in Epileptic EEG Signal using Empirical Mode Decomposition and Support Vector Machine,” *Proc. - 2019 Int. Semin. Intell. Technol. Its Appl. ISITIA 2019*, pp. 314–319, 2019.

- [23] N. Nicolaou and J. Georgiou, “Detection of epileptic electroencephalogram based on Permutation Entropy and Support Vector Machines,” *Expert Syst. Appl.*, vol. 39, no. 1, pp. 202–209, 2012.
- [24] N. H. Farhat, “Photonit neural networks and learning mathines the role of electron-trapping materials,” *IEEE Expert. Syst. their Appl.*, vol. 7, no. 5, pp. 63–72, 1992.
- [25] R. Acharya U., P. S. Bhat, N. Kannathal, A. Rao, and M. L. Choo, “Analysis of cardiac health using fractal dimension and wavelet transformation,” *Itbm-Rbm*, vol. 26, no. 2, pp. 133–139, 2005.
- [26] M. S. J. Solaija, S. Saleem, K. Khurshid, S. A. Hassan, and A. M. Kamboh, “Dynamic mode decomposition based epileptic *seizure* detection from scalp EEG,” *IEEE Access*, vol. 6, no. c, pp. 38683–38692, 2018.
- [27] L. Gautheron, A. Habrard, E. Morvant, and M. Sebban, “Metric learning from imbalanced data,” *Proc. - Int. Conf. Tools with Artif. Intell. ICTAI*, vol. 2019-Novem, no. 9, pp. 923–930, 2019.
- [28] D. M. Hawkins, “The Problem of Overfitting,” *J. Chem. Inf. Comput. Sci.*, vol. 44, no. 1, pp. 1–12, 2004.
- [29] T. N. Alotaiby, S. A. Alshebeili, F. M. Alotaibi, and S. R. Alrshoud, “Epileptic *seizure* prediction using CSP and LDA for scalp EEG signals,” *Comput. Intell. Neurosci.*, vol. 2017, 2017.
- [30] Y. U. Khan, N. Rafiuddin, and O. Farooq, “Automated *seizure* detection in scalp EEG using multiple wavelet scales,” *2012 IEEE Int. Conf. Signal Process. Comput. Control. ISPCC 2012*, pp. 1–5, 2012.