

TABLE I: Performance Comparison Between Models for Different Lookbacks

Model	Lookback	RMSE	MAE	R^2	C.T. (min)
iTransformer	24	0.0041	0.0489	0.9713	0.1
	48	0.0415	0.0323	0.9882	0.3
	96	0.0455	0.0368	0.9863	0.36
TCN	24	0.1050	0.0862	0.9861	3.3
	48	0.1227	0.1006	0.9810	2.3
	96	0.1172	0.0933	0.9827	1.88
Transformer	24	0.0410	0.0331	0.9886	5.05
	48	0.0409	0.0325	0.9883	12.1
	96	0.1285	0.1033	0.9792	19.8

Table I compares the performance of iTransformer, TCN, and Transformer models for lookbacks of 24, 48, and 96. The performance metrics include Root Mean Squared Error ($RMSE$), Mean Absolute Error (MAE), R-squared (R^2), and $C.T.$ representing Computing Time. The bold values indicate the best performance for each metric.




Fig. 6: iTransformer 14-days Actual and Prediction Data.

V. CONCLUSION

We presented sea level forecasting using a novel deep-learning approach called iTransformer with short-term data, specifically six months, utilizing four months for training and the remaining two months for validation and testing. The iTransformer model predicts sea levels over 14 days. As a case study, we used a dataset from Singaraja, Bali, Indonesia. Based on our experiments, it can be concluded that iTransformer achieves the highest prediction accuracy, surpassing both TCN and Transformer models. Additionally, the iTransformer model exhibits lower computational times, requiring only 0.1 minutes for a 24-hour lookback, 0.3 minutes for a 48-hour lookback, and 0.36 minutes for a 96-hour lookback.

We also discovered that a 48-hour lookback period yields high accuracy for the iTransformer model, with an $RMSE$ of 0.0415 and an MAE of 0.0323, both of which are very low. This is complemented by R^2 scores of 0.9882, which indicate strong predictive performance but compared to the other models, iTransformer excels in short computational time. The results of the iTransformer model may serve as a foundational basis for future iterations of the iTransformer as a forecasting model. It is important to note that multivariate datasets are highly recommended, as iTransformer is designed for multivariate analysis, whereas this study employed univariate datasets.

LIMITATIONS AND FUTURE WORKS

Our tests were conducted on univariate and short-term datasets, while iTransformer is designed to excel with long-term multivariate datasets. We suggest further research and improvement are necessary to fully evaluate and implement iTransformer with multivariate features. We sincerely hope our comprehensive studies can benefit future work in this area.

iTransformer possesses various advantages and disadvantages. Nonetheless, it offers extensive opportunities to explore, reproduce, and implement long-term multivariate datasets.

REFERENCES

- [1] IPCC. Chapter 4: Sea level rise and implications for low-lying islands, coasts and communities, March 2024. *Cross-Chapter Box 5 in Chapter 1, 4.2.2, 4.2.2.1, 4.2.2.5, SM 4.1* Non-climatic anthropogenic drivers, including recent and historical demographic and settlement trends and anthropogenic subsidence, have played an important role in increasing low-lying coastal communities' exposure and vulnerability to SLR and extreme sea level (ESL) events (very high confidence). In coastal deltas, for example, these drivers have altered freshwater and sediment availability (high confidence).
- [2] Gary Griggs. Rising seas in california—an update on sea-level rise science. In *World Scientific Encyclopedia of Climate Change: Case Studies of Climate Risk, Action, and Opportunity Volume 3*, pages 105–111. World Scientific, 2021.
- [3] Sida Li, Lintao Liu, Song Cai, and Guocheng Wang. Tidal harmonic analysis and prediction with least-squares estimation and inaction method. *Estuarine, Coastal and Shelf Science*, 220:196–208, 2019.
- [4] F Masri, D Saepudin, and D Adytia. Forecasting of sea level time series using deep learning rnn, lstm, and bilstm, case study in jakarta bay, indonesia. *e-Proceeding of Engineering*, 7(2):8544–8551, 2020.
- [5] Avelisa Yoelma Winona and Didit Adytia. Short term forecasting of sea level by using lstm with limited historical data. In *2020 International Conference on Data Science and Its Applications (ICoDSA)*, pages 1–5, 2020.
- [6] Ridho Nobolino Sabillah and Didit Adytia. Time series forecasting of sea level by using transformer approach, with a case study in pangandaran, indonesia. In *2023 IEEE 8th International Conference for Convergence in Technology (I2CT)*, pages 1–6, 2023.
- [7] Deni Saepudin, Egi Shidqi Rabbani, Dio Navalady, and Didit Adytia. Water level time series forecasting using tcn study case in surabaya. *Jurnal Online Informatika*, 9(1):61–69, 2024.
- [8] Yong Liu, Tengge Hu, Haoran Zhang, Haixu Wu, Shiyu Wang, Lintao Ma, and Mingsheng Long. itransformer: Inverted transformers are effective for time series forecasting. *arXiv preprint arXiv:2310.06625*, 2023.
- [9] Ronald Tulus, Didit Adytia, Nugrahinggil Subasita, and Dede Tarwidi. Sea level prediction by using seasonal autoregressive integrated moving average model, case study in semarang, indonesia. In *2020 8th International Conference on Information and Communication Technology (ICoICT)*, pages 1–5, 2020.
- [10] Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. In I. Guyon, U. Von Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 30. Curran Associates, Inc., 2017.
- [11] Ailing Zeng, Muxi Chen, Lei Zhang, and Qiang Xu. Are transformers effective for time series forecasting?, 2022.